



Creatinase

ORIGIN: RECOMBINANT *E. COLI*

CAT#: CRE-70-2421

EC#: 3.5.3.3

SPECIFICATIONS

Appearance: White to light yellow lyophilizate

Activity: ≥ 9 U/mg lyophilizate

ASSAY PRINCIPLE

Creatinase catalyzes the following reaction:



The appearance of urea is measured spectrophotometrically at 435 nm.

APPLICATION

The enzyme is useful for the determination of creatinine and creatine in clinical analysis.

UNIT DEFINITION

One unit (U) is defined as the amount of enzyme which produces 1 μmol of urea per min at 37°C and pH 7.7 under standard assay conditions.

CHARACTERISTICS

Molecular weight: ca. 80 kDa (gel filtration)

Structure: 2 subunits of 46 kDa (SDS-PAGE)

Michaelis constant: 1.3×10^{-2} M (creatinine)

pH Optimum: 7.0–9.0

pH Stability: 5.0–11.0

Optimum temperature: 40°C

Thermal stability: below 45°C

Stability (liquid form): stable at 37°C for at least two weeks

Stability (powder form): stable at 30°C for at least one month

Inhibitor: Hg^{2+}

Creatinase

Figure -1 pH Optimum




Figure -2 pH Stability




Figure -3 Optimum temperature





Figure -4 Thermal stability



THE AMERICAS

SEKISUI Diagnostics, LLC
 One Wall Street
 Burlington, MA 01803
 Phone: 800 332 1042
 Fax: 800 762 6311

info@sekisui-dx.com
 sekisuidiagnostics.com

INTERNATIONAL

SEKISUI Diagnostics (UK) Limited
 Liphook Way, Allington
 Maidstone, Kent, ME16 0LQ, UK
 Phone: +44 1622 607800
 Fax: +44 1622 607801



Because every result matters™